


# Applications of Fibonacci Numbers

Volume 9

edited by

Fredric T. Howard



KLUWER ACADEMIC PUBLISHERS

# Applications of Fibonacci Numbers

Volume 9

Proceedings of The Tenth International Research Conference  
on Fibonacci Numbers and Their Applications

edited by

Frederic T. Howard

*Wake Forest University,  
Winston-Salem, North Carolina, U.S.A.*



KLUWER ACADEMIC PUBLISHERS  
DORDRECHT / BOSTON / LONDON

# FIBINOMIAL IDENTITIES

Arthur T. Benjamin, Jennifer J. Quinn and Jeremy A. Rouse

Problem A – 6 from the 1990 Putnam exam states:

If  $X$  is a finite set, let  $|X|$  denote the number of elements in  $X$ . Call an ordered pair  $(S, T)$  of subsets of  $\{1, 2, \dots, n\}$  *admissible* if  $s > |T|$  for each  $s \in S$ , and  $t > |S|$  for each  $t \in T$ . How many admissible ordered pairs of subsets of  $\{1, 2, \dots, 10\}$  are there? Prove your answer.

It is no coincidence that the solution, 17711, is the 21<sup>st</sup> Fibonacci number. The number of admissible ordered pairs of subsets of  $\{1, 2, \dots, n\}$  with  $|S| = a$  and  $|T| = b$  is  $\binom{n-b}{a} \binom{n-a}{b}$ . As we shall show, summing over all values of  $a$  and  $b$  leads to

## Identity 1:

$$\sum_{a=0}^n \sum_{b=0}^n \binom{n-b}{a} \binom{n-a}{b} = f_{2n+1}.$$

where  $f_0 = 1$ ,  $f_1 = 1$  and for  $n \geq 2$ ,  $f_n = f_{n-1} + f_{n-2}$ . The published solutions [6, 7] use “convoluted” algebraic methods. Yet the presence of both Fibonacci numbers and binomial coefficients demands a combinatorial explanation. Beginning with our proof of Identity 1, we provide simple, combinatorial arguments for many *fibinomial identities* – identities that combine (generalized) Fibonacci numbers and binomial coefficients.

Fibonacci numbers can be combinatorially interpreted in many ways [10]. The primary tool used in this note will be tilings of  $1 \times n$  boards with tiles of varying lengths. The identities presented are viewed as counting questions, answered in two different ways. To begin with, Identity 1 is easily seen by answering

~~This paper is in final form and no version of it will be submitted for publication elsewhere.~~

**Question:** How many ways can a board of length  $2n + 1$  be tiled using (length 1) squares and (length 2) dominoes?

**Answer 1:** A length  $m$  board can be tiled in  $f_m$  ways, which can be seen by conditioning on whether the last tile is a square or a domino. Consequently, a board of length  $2n + 1$  can be tiled  $f_{2n+1}$  ways.

**Answer 2:** Condition on the number of dominoes on each side of the *median square*.

Any tiling of a  $(2n + 1)$ -board must contain an odd number of squares. Thus one square, which we call the median square, contains an equal number of squares to the left and right of it. For example, the 13-tiling in Figure 1 has 5 squares. The median square, the third square, is located in cell 9.

How many tilings contain exactly  $a$  dominoes to the left of the median square and exactly  $b$  dominoes to the right of the median square? Such a tiling has  $(a + b)$  dominoes and therefore  $(2n + 1) - 2(a + b)$  squares. Hence the median square has  $n - a - b$  squares on each side of it. Since the left side has  $(n - a - b) + a = n - b$  tiles, of which  $a$  are dominoes, there are  $\binom{n-b}{a}$  ways to tile to the left of the median square. Similarly, there are  $\binom{n-b}{b}$  ways to tile to

the right of the median square. Hence there are  $\binom{n-b}{a} \binom{n-b}{b}$  tilings altogether.

Varying  $a$  and  $b$  over all feasible values, we obtain the total number of  $(2n + 1)$  tilings as  $\sum_{a=0}^n \sum_{b=0}^n \binom{n-b}{a} \binom{n-b}{b}$ .



Figure 1: Every square-domino tiling of odd length must have a median square. The 13-tiling above has 3 dominoes left of the median square and 1 domino to the right of the median square. The number of such tilings is  $\binom{5}{3} \binom{3}{1}$ .

We can extend this identity by utilizing the 3-bonacci numbers, defined by  $\theta_n = 0$  for  $n < 0$ ,  $\theta_0 = 1$  and for  $n \geq 1$ ,  $\theta_n = \theta_{n-1} + \theta_{n-3}$ .

**Identity 2:**

$$\sum_{a=0}^n \sum_{b=0}^n \sum_{c=0}^n \binom{n-b-c}{a} \binom{n-a-c}{b} \binom{n-a-b}{c} = \theta_{3n+2}.$$

**Question:** How many ways can a board of length  $3n + 2$  be tiled using squares and trominoes?

**Answer 1:** It is easy to see that  $\theta_n$  counts the number of ways to tile a board of length  $n$  with squares and (length 3) trominoes. Hence there are  $\theta_{3n+2}$  such tilings of a board of length  $3n + 2$ .

**Answer 2:** The number of squares in any tiling of a  $(3n + 2)$ -board must be 2 greater than a multiple of 3. Hence there will exist two *goalpost* squares, say located at cells  $x$  and  $y$ , such that there are an equal number of squares to the left of  $x$ , between  $x$  and  $y$ , and to the right of  $y$ . We condition on the number of trominoes in the three regions defined by the goalposts. If the number of trominoes in each region is, from left to right,  $a, b, c$ , then there are a total of  $a + b + c$  trominoes and  $(3n + 2) - 3(a + b + c)$  squares, including the two goalpost squares.

Thus each region has  $n - (a + b + c)$  squares. The leftmost region has  $n - b - c$  tiles,  $a$  of which are trominoes, and there are  $\binom{n-b-c}{a}$  ways to arrange them. Likewise the tiles of the second

and third region can be arranged  $\binom{n-a-c}{b}$  ways and  $\binom{n-a-b}{c}$  ways, respectively.

As  $a$ ,  $b$ , and  $c$  vary, the total number of  $(3n + 2)$ -tilings is the left side of our identity.

Applying the same logic, using only tiles of length 1 and  $k$ , we immediately obtain the following  $k$ -bonacci generalization.

**Identity 3:** Let  $\kappa_n$  be the  $k$ -bonacci number defined by  $\kappa_n = 0$  for  $n < 0$ ,  $\kappa_0 = 1$ , and for  $n \geq 1$ ,  $\kappa_n = \kappa_{n-1} + \kappa_{n-k}$ . Then for  $n \geq 0$ ,  $\kappa_{kn+(k-1)}$  equals

$$\sum_{a_1=0}^n \sum_{a_2=0}^n \cdots \sum_{a_k=0}^n \binom{n - (a_2 + a_3 + \cdots + a_k)}{a_1} \binom{n - (a_1 + a_3 + \cdots + a_k)}{a_2} \cdots \binom{n - (a_1 + a_2 + \cdots + a_{k-1})}{a_k} \quad (1)$$

Another generalization of Fibonacci numbers are the  $k^{\text{th}}$  order Fibonacci numbers defined by  $g_n = 0$  for  $n < 0$ ,  $g_0 = 1$ , and for  $n \geq 1$ ,  $g_n = g_{n-1} + g_{n-2} + \cdots + g_{n-k}$ . The next identity is proved by non-trivial algebraic methods in [8] and [9], but when viewed combinatorially, as done in [2], it is practically obvious.

**Identity 4:** For all  $n \geq 0$ ,

$$\sum_{n_1} \sum_{n_2} \cdots \sum_{n_k} \binom{n_1 + n_2 + \cdots + n_k}{n_1, n_2, \dots, n_k} = g_n,$$

the  $k^{\text{th}}$  order Fibonacci number, where the summation is over all non negative integers  $n_1, n_2, \dots, n_k$  such that  $n_1 + 2n_2 + \cdots + kn_k = n$ .

**Question:** In how many ways can we tile a board of length  $n$  using tiles with lengths at most  $k$ ?

**Answer 1:** By its definition, it is combinatorially clear that  $g_n$  counts this quantity.

**Answer 2:** Condition on the number of tiles of each length. If for  $1 \leq i \leq k$  there are  $n_i$  tiles of length  $i$ , then we must have  $n_1 + 2n_2 + \cdots + kn_k = n$ . The number of ways to permute these tiles is given by the multinomial coefficient  $\binom{n_1+n_2+\dots+n_k}{n_1, n_2, \dots, n_k}$ .

More colorfully, for nonnegative integers  $c_1, \dots, c_k$  we define the generalized  $k^{\text{th}}$  order Fibonacci number by  $h_n = 0$  for  $n < 0$ ,  $h_0 = 1$ , and for  $n \geq 1$ ,  $h_n = c_1 \cdot h_{n-1} + c_2 \cdot h_{n-2} + \cdots + c_k \cdot h_{n-k}$ . It is easy to see that  $h_n$  counts the number of ways to tile a board of length  $n$  with colored tiles of length at most  $k$ , where for  $1 \leq i \leq k$ , a tile of length  $i$  may be assigned any one of  $c_i$  colors. The previous identity and argument immediately generalizes to the following identity.

**Identity 5:** For all  $n \geq 0$ ,

$$\sum_{n_1} \sum_{n_2} \cdots \sum_{n_k} \binom{n_1 + n_2 + \cdots + n_k}{n_1, n_2, \dots, n_k} c_1^{n_1} c_2^{n_2} \cdots c_k^{n_k} = h_n,$$

the generalized  $k^{\text{th}}$  order Fibonacci number, where the summation is over all non-negative integers  $n_1, n_2, \dots, n_k$  such that  $n_1 + 2n_2 + \cdots + kn_k = n$ .

Our tiling approach also succeeds in proving even more complex fibinomial identities.

**Identity 6:**

$$\sum_{a_1=0}^n \sum_{a_2=0}^n \cdots \sum_{a_k=0}^n \binom{n}{a_1} \binom{n-a_1}{a_2} \cdots \binom{n-a_{k-1}}{a_k} = f_{k+1}^n.$$

**Question:** In how many ways can we simultaneously tile  $n$  distinguishable boards of length  $k + 1$  with squares and dominoes?

**Answer 1:** Since each board can be tiled  $f_{k+1}$  ways, there are  $f_{k+1}^n$  such tilings.

**Answer 2:** Condition on the number of dominoes covering each consecutive pair of cells. We claim there are  $\binom{n}{a_1} \binom{n-a_1}{a_2} \cdots \binom{n-a_{k-1}}{a_k}$  ways to create  $n$  tilings of length  $k + 1$  where  $a_1$  of them begin with dominoes,  $a_2$  have dominoes covering cells 2 and 3, and generally for  $1 \leq i \leq k$ ,  $a_i$  of them have dominoes covering cells  $i$  and  $i + 1$ . To see this, notice there are  $\binom{n}{a_1}$  ways to decide which of the  $n$  tilings begin with a domino (the rest begin with a square). Once these are selected, then among those  $n - a_1$  tilings that do not begin with a domino there are  $\binom{n-a_1}{a_2}$  ways to determine which of those will have a domino in cells 2 and 3. (The other  $n - a_1 - a_2$  tilings will have a square in cell 2.) Continuing in this fashion, we see that once the tilings with dominoes covering cells  $i - 1$  and  $i$  are determined, there are  $\binom{n-a_{i-1}}{a_i}$  ways to determine which tilings have dominoes covering cells  $i$  and  $i + 1$ .

More generally, by tiling  $n$  distinguishable boards of length  $k + 1$  with squares and dominoes where the first  $c$  of them must begin with a square, the same reasoning establishes:

**Identity 7:** For  $0 \leq c \leq n$ ,

$$\sum_{a_1=0}^n \sum_{a_2=0}^n \cdots \sum_{a_k=0}^n \binom{n-c}{a_1} \binom{n-a_1}{a_2} \cdots \binom{n-a_{k-1}}{a_k} = f_k^c f_{k+1}^{n-c}.$$

Identities 6 and 7 can be extended to *Gibonacci* numbers,  $G_n$ , defined by initial conditions  $G_0, G_1$ , and the Fibonacci recurrence  $G_n = G_{n-1} + G_{n-2}$  for  $n \geq 2$ . For non-negative integers  $G_0$  and  $G_1$ ,  $G_n$  can be combinatorially defined [3] as the number of ways to tile a length  $n$  board with squares and dominoes subject to the *initial conditions* that the first tile is given a *phase*, where there are  $G_0$  choices for the phase of a domino and  $G_1$  choices for the phase of a square. The extension of Identity 6 to Gibonacci numbers is

**Identity 8:**

$$\sum_{a_1=0}^n \sum_{a_2=0}^n \dots \sum_{a_k=0}^n \binom{n}{a_1} \binom{n-a_1}{a_2} \dots \binom{n-a_{k-1}}{a_k} G_0^{a_1} G_1^{n-a_1} = G_{k+1}^n.$$

**Question:** In how many ways can we simultaneously tile  $n$  distinguishable boards of length  $k + 1$  with squares and dominoes, where an initial domino is assigned one of  $G_0$  phases and an initial square is assigned one of  $G_1$  phases?

**Answer 1:** Since each board can be tiled  $G_{k+1}$  ways, there are  $G_{k+1}^n$  such tilings.

**Answer 2:** As in the proof of Identity 6, we condition on the number of dominoes covering each consecutive pair of cells. The difference here is that initially choosing  $a_1$  of the tilings to begin with a domino contributes  $\binom{n}{a_1} G_0^{a_1} G_1^{n-a_1}$  to the product since we must assign domino phases to  $a_1$  of the boards and square phases to the remaining  $n - a_1$  boards.

As an immediate corollary, we have

**Identity 9:**

$$\sum_{a_1=0}^n \sum_{a_2=0}^n \dots \sum_{a_k=0}^n \binom{n}{a_1} \binom{n-a_1}{a_2} \dots \binom{n-a_{k-1}}{a_k} 2^{a_1} = L_{k+1}^n,$$

where  $L_n$  is the  $n^{th}$  Lucas number.

Likewise, if we require that the first  $c$  of the  $n$  boards begin with phased squares, then the same reasoning establishes the following Gibonacci extension of Identity 7.

**Identity 10:** For  $0 \leq c \leq n$ ,

$$\sum_{a_1=0}^n \sum_{a_2=0}^n \dots \sum_{a_k=0}^n \binom{n-c}{a_1} \binom{n-a_1}{a_2} \dots \binom{n-a_{k-1}}{a_k} G_0^{a_1} G_1^{n-a_1} = G_1^c f_k^c G_{k+1}^{n-c}$$

As with Identity 5, these identities can be further generalized by colorizing them. For more combinatorial proofs of fibinomial identities, see [4].

We end with an open question. The *Fibonomial Numbers* are defined like binomial coefficients with  $F$ 's on top. That is, Fibinomial  $\binom{n}{k}_F = \frac{F_n F_{n-1} \dots F_{n-k+1}}{F_1 F_2 \dots F_k}$ , where  $F_n = f_{n-1}$  is the traditional  $n^{th}$  Fibonacci number. Amazingly  $\binom{n}{k}_F$  is always an integer [1]. We challenge the reader to find a combinatorial proof of this fact.

**ACKNOWLEDGMENT**

We thank the referee for many helpful suggestions. This research was supported by The Reed Institute for Decision Science and the Beckman Research Foundation.

REFERENCES

- [1] Alexanderson, G.L. and Klosinski, L.F. "A Fibonacci Analogue of Gaussian Binomial coefficients." *The Fibonacci Quarterly*, Vol. 12 (1974): pp. 129-132.
- [2] Benjamin, A.T., Hanusa, C.R.H. and Su, F.E. "Linear Recurrences Through Tilings and Markov Chains." *Utilitas Mathematica*, Vol. 64 (2003): pp. 3-17.
- [3] Benjamin, A.T., Quinn, J.J. and Su, F.E. "Generalized Fibonacci Identities through Phased Tilings." *The Fibonacci Quarterly*, Vol. 38.3 (2000): pp. 282-288.
- [4] Benjamin, A.T. and Rouse, J.A. "Recounting Binomial Fibonacci Identities." Applications of Fibonacci Numbers. Volume 10. Edited by Fredric T. Howard. Kluwer Academic Publishers, Dordrecht, The Netherlands, 2003: pp. 25-28.
- [5] Carlitz, L. "The Characteristic Polynomial of a Certain Matrix of Binomial Coefficients." *The Fibonacci Quarterly*, Vol. 3.2 (1965): pp. 81-89.
- [6] Klosinski, L.F. Alexanderson, G.L. and Larson, L.C. "The Fifty-First William Lowell Putnam Mathematical Competition." *American Mathematical Monthly*, Vol. 98.8 (1991): pp. 719-727.
- [7] Larson, L. "The Fifty-First William Lowell Putnam Mathematical Competition." *Mathematics Magazine*, Vol. 64.2 (1991): pp. 141-144.
- [8] Philippou, A.N. "A Note on the Fibonacci Sequence of Order  $K$  and the Multinomial Coefficients." *The Fibonacci Quarterly*, Vol. 21.2 (1983): pp. 82-86.
- [9] Philippou, A.N. and Muwafi, A.A. "Waiting for the  $K$ -th Consecutive Success and the Fibonacci Sequence of Order  $K$ ." *The Fibonacci Quarterly*, Vol. 20.1 (1982): pp. 28-32.
- [10] Stanley, R.P. Enumerative Combinatorics. Volume 1 *Wadsworth and Brooks/Cole* Chapter 1 problem 14a-i.

AMS Classification Numbers: 05A19, 11B39