



## What's Best?

Arthur T. Benjamin; Matthew T. Fluet

*The American Mathematical Monthly*, Vol. 107, No. 6 (Jun. - Jul., 2000), 560-562.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28200006%2F07%29107%3A6%3C560%3AWB%3E2.0.CO%3B2-K>

*The American Mathematical Monthly* is currently published by Mathematical Association of America.

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/maa.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

---

JSTOR is an independent not-for-profit organization dedicated to creating and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

details to the reader. The first Hankel determinant is classical, the second is due to Radoux [3].

**Theorem 4.** *Let  $D(n)$  denote the number of derangements of an  $n$ -element set, that is, the number of permutations without fixed points. The Hankel determinants of order  $n + 1$  of the matrices  $[(i + j)!]$  and  $[D(i + j)]$  are given by*

$$\det((i + j)!)_{0 \leq i, j \leq n} = \det(D(i + j))_{0 \leq i, j \leq n} = \left( \prod_{i=0}^n i! \right)^2.$$

**ACKNOWLEDGMENTS.** The author thanks Margaret Readdy for reading an earlier draft of this note. This work was done when the author was a member at the Institute for Advanced Study where he was supported by National Science Foundation, DMS 97-29992, and NEC Research Institute, Inc.

#### REFERENCES

1. M. Gardner, *Fractal Music, Hypercards and more: Mathematical Recreations from Scientific American*, W. H. Freeman and Company, New York, 1992.
2. C. Radoux, Calcul effectif de certains déterminants de Hankel, *Bull. Soc. Math. Belg. Sér. B* **31** (1979) 49–55
3. C. Radoux, Déterminant de Hankel construit sur des polynomes liés aux nombres de dérangements, *European J. Combin.* **12** (1991) 327–329.
4. C. Radoux, Déterminants de Hankel et théorème de Sylvester, in *Séminaire Lotharingien de Combinatoire (Saint-Nabor, 1992)*, *Publ. Inst. Rech. Math. Av.*, 498, Univ. Louis Pasteur, Strasbourg, 1992, pp. 115–122.
5. C. Radoux, Addition formulas for polynomials built on classical combinatorial sequences, to appear in *J. Comput. Appl. Math.*
6. G.-C. Rota, *Finite Operator Calculus*, Academic Press, Inc., New York, 1975.
7. G.-C. Rota, D. Kahaner, and A. Odlyzko, On the foundations of combinatorial theory. VIII. Finite operator calculus, *J. Math. Anal. Appl.* **42** (1973) 684–760.


Royal Institute of Technology, S-100 44 Stockholm, Sweden  
 jrge@math.kth.se

## What's Best?

**Arthur T. Benjamin and Matthew T. Fluet**

Consider the following game. You are given 9 tokens and a weighted coin with heads probability  $2/3$ . You allocate the tokens so that some are assigned to heads and the rest assigned to tails. (See Figure 1 for two examples.) Next the coin is flipped, and if heads (tails) is flipped, then one token assigned to heads (tails) is removed. The coin is flipped until all tokens are removed. The challenge is to determine how to allocate the tokens so that they are removed as quickly as possible. Which of the allocations in Figure 1 is best?


Surprisingly, the answer depends on what you mean by best. Allocation  $A = (6, 3)$  is attractive since it is proportional to the probabilities. Nonetheless, if you wish to minimize the average number of flips needed to remove all tokens, then  $B = (7, 2)$  turns out to be best. On the other hand, when allocations  $A$  and  $B$  compete against each other using the same coin, you should bet on  $A$  to finish



**Figure 1.** Using a coin with heads probability  $2/3$ , which card is cleared faster?

before  $B$ . In fact, when competing with any other allocation  $C$  of 9 tokens,  $A$  is favored to finish before  $C$ .

Why does  $A$  beat  $B$  more often than not? Let  $X$  denote the number of heads obtained by the coin (whose outcomes apply to both players) after the first 9 flips. If  $X \leq 6$ , then  $A$  must eventually win since both players have removed all of their tail tokens, and  $A$  has one fewer head token remaining. By the histogram of probabilities in Figure 2, we see  $P(X \leq 6) = 0.6228$ . The same argument shows that  $A$  is also favored to beat any allocation of 9 tokens with more than 7 heads. Likewise, allocation  $A$  will have a winning record against allocations of 9 tokens with fewer than 6 heads since  $A$  will win all contests whenever  $X \geq 6$ , which has probability 0.6503.



**Figure 2.** The median of the  $\text{BINOMIAL}(9, \frac{2}{3})$  distribution.

What makes  $(6, 3)$  best in competition is that 6 is the *median* of the binomial distribution with parameters 9 and  $2/3$ . In general, we have the following:

**Theorem 1.** Using a coin with heads probability  $p$ , the best (heads, tails) allocation of  $t$  tokens is  $(m, t - m)$ , where  $m$  is the median of the  $\text{BINOMIAL}(t, p)$  distribution. It is best in the sense that it is favored to finish sooner than any other allocation of  $t$  tokens.

Thus when faced with an opponent, the optimal number of heads is the .50 percentile of the  $\text{BINOMIAL}(t, p)$  distribution. However when playing alone, we have

**Theorem 2.** Using a coin with heads probability  $p$ , the best (heads, tails) allocation of  $t$  tokens is  $(m, t - m)$ , where  $m$  is the  $p$ -th percentile of the  $\text{BINOMIAL}(t, p)$  distribution. It is best in the sense that it has the smallest expected number of flips among all allocations of  $t$  tokens.

Theorem 2 asserts that the best allocation of 9 tokens is (7, 2) since  $P(X \leq 6) = 0.6228 \leq 2/3 \leq 0.8569 = P(X \leq 7)$ . A recursive calculation shows that on average, allocation (7, 2) uses 11.3 flips whereas (6, 3) uses 11.5 flips to clear.

To prove Theorem 2, let  $E(n)$  denote the average clearing time of allocation  $(n, t - n)$ . Although a formula for  $E(n)$  can be computed explicitly [2], it is not necessary for our proof. Instead, we show that  $E(n + 1) \geq E(n)$  if and only if  $n$  is greater than or equal to the  $p$ -th percentile of the  $\text{BINOMIAL}(t, p)$  distribution.

We compute  $E(n + 1) - E(n)$  by conditioning on the  $\text{BINOMIAL}(t, p)$  random variable  $X$ , the number of heads obtained in the first  $t$  flips of the coin. If  $X \leq n$ , then both allocations will have their “tails cleaned”, and the resulting allocation  $(n + 1 - X, 0)$  will need one more head than  $(n - X, 0)$ , which takes on average  $1/p$  flips longer. On the other hand, if  $X > n$ , then the resulting allocation  $(0, X - n - 1)$  will finish on average  $1/(1 - p)$  flips before  $(0, X - n)$ . Thus

$$\begin{aligned} E(n + 1) - E(n) &= \frac{1}{p} P(X \leq n) - \frac{1}{1 - p} [1 - P(X \leq n)] \\ &= \frac{P(X \leq n) - p}{p(1 - p)}, \end{aligned}$$

and the result follows.

In [3] it is shown that the median of the  $\text{BINOMIAL}(t, p)$  distribution can not differ from the mean  $tp$  by more than  $\ln 2$ . In particular, when  $tp$  is an integer, the mean, median, and mode are all equal. When  $t$  is large, the central limit theorem asserts that the  $p$ -th percentile is approximately  $tp + z_p \sqrt{tp(1 - p)}$ , where  $z_p$  is the  $p$ -th percentile of the standard normal distribution. Thus, no matter what definition of “best” you prefer, in the long run, the best allocation will be approximately  $(tp, t(1 - p))$ .

Suppose the game is played with a biased *die* with  $s$  sides instead of a biased coin? Although some necessary conditions are given in [2], there is no known closed form for the allocations with minimum average clearing time. To make matters more interesting [1], many values of  $t$  and  $(p_1, p_2, \dots, p_s)$  produce non-transitive situations in which allocations  $A$ ,  $B$ , and  $C$  are favored against all *other* allocations, but  $A$  is favored against  $B$ ,  $B$  is favored against  $C$ , and  $C$  is favored against  $A$ . What’s best in this situation? Ask your opponent to allocate the tokens first!

**ACKNOWLEDGMENT.** We thank Janet Myhre and The Reed Institute of Applied Mathematics for supporting this work.

#### REFERENCES

1. A. T. Benjamin and M. T. Fluet, The Best Way to Knock ’m Down, *The UMAP Journal* 20.1 (1999) 11–20.
2. M. T. Fluet, Searching for Optimal Strategies in Knock ’m Down, *Senior Thesis*, Mathematics Department, Harvey Mudd College, 1999.
3. K. Hamza, The smallest uniform upper bound on the distance between the mean and the median of the binomial and Poisson distributions, *Statist. Probab. Lett.* 23 (1995) 21–25.

Harvey Mudd College, Claremont, CA 91711  
 benjamin@hmc.edu  
 fluet@cs.cornell.edu